GUIDELINES OPENING UP AMERICA AGAIN





Proposed State or Regional Gating Criteria

(Satisfy Before Proceeding to Phased Opening)

SYMPTOMS

Downward trajectory of influenza-like illnesses (ILI) reported within a 14-day period

AND

Downward trajectory of covid-like syndromic cases reported within a 14-day period

CASES

Downward trajectory of documented cases within a 14-day period

OR

Downward trajectory of positive tests as a percent of total tests within a 14-day period (flat or increasing volume of tests)

*State and local officials may need to tailor the application of these criteria to local circumstances (e.g., metropolitan areas that have suffered severe COVID outbreaks, rural and suburban areas where outbreaks have not occurred or have been mild). Additionally, where appropriate, Governors should work on a regional basis to satisfy these criteria and to progress through the phases outlined below.

HOSPITALS

Treat all patients without crisis care

AND

Robust testing program in place for at-risk healthcare workers, including emerging antibody testing



Core State Preparedness Responsibilities

TESTING & CONTACT TRACING HEALTHCARE SYSTEM CAPACITY

✓ Ability to quickly set up safe and efficient ✓ Ability to quickly and independently supply sufficient Personal Protective screening and testing sites for symptomatic Equipment and critical medical equipment to handle dramatic surge in need individuals and trace contacts of COVID+ ✓ Ability to surge ICU capacity results **PLANS** Ability to test Syndromic/ILI-indicated persons \checkmark ✓ Protect the health and safety of workers in critical industries for COVID and trace contacts of COVID+ results ✓ Protect the health and safety of those living and working in high-risk facilities (e.g., ✓ Ensure sentinel surveillance sites are senior care facilities) screening for asymptomatic cases and ✓ Protect employees and users of mass transit contacts for COVID+ results are traced (sites ✓ Advise citizens regarding protocols for social distancing and face coverings operate at locations that serve older Monitor conditions and immediately take steps to limit and mitigate any rebounds or individuals, lower-income Americans, racial outbreaks by restarting a phase or returning to an earlier phase, depending on severity minorities, and Native Americans)







Proposed Phased Approach

> IMPLEMENTABLE ON STATEWIDE OR COUNTY-**BY-COUNTY** BASIS AT GOVERNORS' DISCRETION

BASED ON UP-TO-DATE DATA AND READINESS

MITIGATES RISK OF RESURGENCE

PROTECTS THE MOST VULNERABLE

Guidelines for All Phases: Individuals


CONTINUE TO PRACTICE GOOD HYGIENE

- ✓ Wash your hands with soap and water or use hand sanitizer, especially after touching frequently used items or surfaces.
- ✓ Avoid touching your face.
- ✓ Sneeze or cough into a tissue, or the inside of your elbow.
- Disinfect frequently used items and surfaces as much as possible. \checkmark
- ✓ Strongly consider using face coverings while in public, and particularly when using mass transit.

PEOPLE WHO FEEL SICK SHOULD STAY HOME

- ✓ Do not go to work or school.
- ✓ Contact and follow the advice of your medical provider.

Continue to adhere to State and local guidance as well as complementary CDC guidance, particularly with respect to face coverings.



Guidelines for All Phases: Employers

Develop and implement appropriate policies, in accordance with Federal, State, and local regulations and guidance, and informed by industry best practices, regarding:

- ✓ Social distancing and protective equipment
- ✓ Temperature checks
- ✓ Testing, isolating, and contact tracing
- ✓ Sanitation
- ✓ Use and disinfection of common and high-traffic areas
- ✓ Business travel

work until cleared by a medical provider.

Develop and implement policies and procedures for workforce contact tracing following employee COVID+ test.

Monitor workforce for indicative symptoms. Do not allow symptomatic people to physically return to















Phase One

FOR STATES AND REGIONS THAT SATISFY THE GATING CRITERIA



Phase One INDIVIDUALS

ALL VULNERABLE INDIVIDUALS* should continue to shelter in place. Members of households with vulnerable residents should be aware that by returning to work or other environments where distancing is not practical, they could carry the virus back home. Precautions should be taken to isolate from vulnerable residents.

All individuals, **WHEN IN PUBLIC** (e.g., parks, outdoor recreation areas, shopping areas), should maximize physical distance from others. Social settings of more than 10 people, where appropriate distancing may not be practical, should be avoided unless precautionary measures are observed.

Avoid **SOCIALIZING** in groups of more than 10 people in circumstances that do not readily allow for appropriate physical distancing (e.g., receptions, trade shows)

MINIMIZE NON-ESSENTIAL TRAVEL and adhere to CDC guidelines regarding isolation following travel.

EACH PHASE OF THIS GUIDANCE ADDRESSES THOSE ASPECTS OF DAILY LIFE FOR WHICH RESTRICTIONS REMAIN APPROPRIATE DUE TO COVID.

*See Appendix 1 for Definition of Vulnerable Individuals









Continue to **ENCOURAGE TELEWORK**, whenever possible and feasible with business operations.

Phase One **EMPLOYERS**

Minimize **NON-ESSENTIAL TRAVEL** and adhere to CDC guidelines regarding isolation following travel.

Strongly consider **SPECIAL ACCOMMODATIONS** for personnel who are members of a **VULNERABLE POPULATION**.

EACH PHASE OF THIS GUIDANCE ADDRESSES THOSE ASPECTS OF DAILY LIFE FOR WHICH RESTRICTIONS REMAIN APPROPRIATE DUE TO COVID.

If possible, **RETURN TO WORK IN PHASES.**

Close **COMMON AREAS** where personnel are likely to congregate and interact, or enforce strict social distancing protocols.





Phase One SPECIFIC TYPES OF **EMPLOYERS**

SCHOOLS AND ORGANIZED YOUTH ACTIVITIES (e.g., daycare, camp) that are currently closed should remain closed.

VISITS TO SENIOR LIVING FACILITIES AND HOSPITALS should be prohibited. Those who do interact with residents and patients must adhere to strict protocols regarding hygiene.

LARGE VENUES (e.g., sit-down dining, movie theaters, sporting venues, places of worship) can operate under strict physical distancing protocols.

ELECTIVE SURGERIES can resume, as clinically appropriate, on an outpatient basis at facilities that adhere to CMS guidelines.

GYMS can open if they adhere to strict physical distancing and sanitation protocols.

BARS should remain closed.

EACH PHASE OF THIS GUIDANCE ADDRESSES THOSE ASPECTS OF DAILY LIFE FOR WHICH RESTRICTIONS REMAIN APPROPRIATE DUE TO COVID.









Phase Two

FOR STATES AND REGIONS WITH NO EVIDENCE OF A REBOUND AND THAT SATISFY THE GATING CRITERIA A SECOND TIME





Phase Two INDIVIDUALS

ALL VULNERABLE INDIVIDUALS should continue to shelter in place. Members of households with vulnerable residents should be aware that by returning to work or other environments where distancing is not practical, they could carry the virus back home. Precautions should be taken to isolate from vulnerable residents.

All individuals, **WHEN IN PUBLIC** (e.g., parks, outdoor recreation areas, shopping areas), should maximize physical distance from others. Social settings of more than 50 people, where appropriate distancing may not be practical, should be avoided unless precautionary measures are observed.

NON-ESSENTIAL TRAVEL can resume.

EACH PHASE OF THIS GUIDANCE ADDRESSES THOSE ASPECTS OF DAILY LIFE FOR WHICH RESTRICTIONS REMAIN APPROPRIATE DUE TO COVID.









Phase Two Employers

Continue to **ENCOURAGE TELEWORK**, whenever possible and feasible with business operations.

Close **COMMON AREAS** where personnel are likely to congregate and interact, or enforce moderate social distancing protocols.

NON-ESSENTIAL TRAVEL can resume.

Strongly consider **SPECIAL ACCOMMODATIONS** for personnel who are members of a **VULNERABLE POPULATION.**

EACH PHASE OF THIS GUIDANCE ADDRESSES THOSE ASPECTS OF DAILY LIFE FOR WHICH RESTRICTIONS REMAIN APPROPRIATE DUE TO COVID.





Phase Two SPECIFIC TYPES OF **EMPLOYERS**

VISITS TO SENIOR CARE FACILITIES AND HOSPITALS should be prohibited. Those who do interact with residents and patients must adhere to strict protocols regarding hygiene.

LARGE VENUES (e.g., sit-down dining, movie theaters, sporting venues, places of worship) can operate under moderate physical distancing protocols.

ELECTIVE SURGERIES can resume, as clinically appropriate, on an outpatient and in-patient basis at facilities that adhere to CMS guidelines.

protocols.

BARS may operate with diminished standing-room occupancy, where applicable and appropriate.

EACH PHASE OF THIS GUIDANCE ADDRESSES THOSE ASPECTS OF DAILY LIFE FOR WHICH RESTRICTIONS REMAIN APPROPRIATE DUE TO COVID.

SCHOOLS AND ORGANIZED YOUTH ACTIVITIES (e.g., daycare, camp) can reopen.

GYMS can remain open if they adhere to strict physical distancing and sanitation



















Phase Three

FOR STATES AND REGIONS WITH NO EVIDENCE OF A REBOUND AND THAT SATISFY THE GATING CRITERIA A THIRD TIME





Phase Three INDIVIDUALS

VULNERABLE INDIVIDUALS can resume public interactions, but should practice physical distancing, minimizing exposure to social settings where distancing may not be practical, unless precautionary measures are observed.

LOW-RISK POPULATIONS should consider minimizing time spent in crowded environments.

Phase Three EMPLOYERS

EACH PHASE OF THIS GUIDANCE ADDRESSES THOSE ASPECTS OF DAILY LIFE FOR WHICH RESTRICTIONS REMAIN APPROPRIATE DUE TO COVID.

Resume UNRESTRICTED STAFFING of worksites.





Phase Three SPECIFIC TYPES OF EMPLOYERS

LARGE VENUES (e.g., sit-down dining, movie theaters, sporting venues, places of worship) can operate under limited physical distancing protocols.

GYMS can remain open if they adhere to standard sanitation protocols.

BARS may operate with increased standing room occupancy, where applicable.

VISITS TO SENIOR CARE FACILITIES AND HOSPITALS can resume. Those who interact with residents and patients must be diligent regarding hygiene.





Appendix Vulnerable Individuals

1. Elderly individuals.

2. Individuals with serious underlying health conditions, including high blood pressure, chronic lung disease, diabetes, obesity, asthma, and those whose immune system is compromised such as by chemotherapy for cancer and other conditions requiring such therapy.